



## **DRAFT EAST AFRICAN STANDARD**

---

**Liquid soaps and detergent for household use — Specification**

**EAST AFRICAN COMMUNITY**

---



### Copyright notice

This EAC document is copyright-protected by EAC. While the reproduction of this document by participants in the EAC standards development process is permitted without prior permission from EAC, neither this document nor any extract from it may be reproduced, stored or transmitted in any form for any other purpose without prior written permission from EAC.

Requests for permission to reproduce this document for the purpose of selling it should be addressed as shown below or to EAC's member body in the country of the requester:

© East African Community 2025 — All rights reserved  
East African Community  
P.O. Box 1096,  
Arusha  
Tanzania  
Tel: + 255 27 2162100  
Fax: + 255 27 2162190  
E-mail: [eac@eachq.org](mailto:eac@eachq.org)  
Web: [www.eac-quality.net](http://www.eac-quality.net)

Reproduction for sales purposes may be subject to royalty payments or a licensing agreement. Violators may be prosecuted.

FOR PUBLIC COMMENT ONLY

## Foreword

Development of the East African Standards has been necessitated by the need for harmonizing requirements governing quality of products and services in the East African Community. It is envisaged that through harmonized standardization, trade barriers that are encountered when goods and services are exchanged within the Community will be removed.

The Community has established an East African Standards Committee (EASC) mandated to develop and issue East African Standards (EAS). The Committee is composed of representatives of the National Standards Bodies in Partner States, together with the representatives from the public and private sector organizations in the community.

East African Standards are developed through Technical Committees that are representative of key stakeholders including government, academia, consumer groups, private sector and other interested parties. Draft East African Standards are circulated to stakeholders through the National Standards Bodies in the Partner States. The comments received are discussed and incorporated before finalization of standards, in accordance with the Principles and procedures for development of East African Standards.

East African Standards are subject to review, to keep pace with technological advances. Users of the East African Standards are therefore expected to ensure that they always have the latest versions of the standards they are implementing.

The committee responsible for this document is Technical Committee EASC/TC 074, *Surface Active agents*.

Attention is drawn to the possibility that some of the elements of this document may be subject of patent rights. EAC shall not be held responsible for identifying any or all such patent rights.

This fourth edition cancels and replaces the third edition (EAS 383:2021), which has been technically revised.

## Introduction

This specification shall apply to all categories of organic and inorganic cleaning agents, including both soap and detergent-based formulations. These include anionic, non-ionic, cationic, and amphoteric surfactants, whether utilized individually or in combination.

Branched alky aryl sulphonates are non-biodegradable. The use of these starting materials is strongly discouraged for environmental conservation. However, when these starting materials are used, they shall be declared by the manufacturer.

FOR PUBLIC COMMENT ONLY

## Liquid soap and detergent for household use — Specification

### 1 Scope

This Draft East African Standard specifies the requirements, sampling and test methods for liquid soap and detergent for household use. This standard does not cover soaps and detergents for laundry use, liquid bathing soap and liquid hand wash (synthetic and combined).

### 2 Normative references

The following documents are referred to in the text in such a way that some or all of their content constitutes requirements of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EAS 377 (all parts), *Cosmetics and cosmetic products*

ISO 2271, *Surface-active agents — Determination of anionic active matter by manual or mechanical direct two-phase titration procedure.*

EAS 814, *Determination of biodegradability of surfactants — Test method*

ISO 862, *Surface active agents — Vocabulary*

ISO 2870, *Surface active agents — Detergents — Determination of anionic-active matter, hydrolysable and non-hydrolysable under acid conditions*

ISO 2871-1, *Surface active agents — Detergents — Determination of cationic-active matter content — Part 1: High-molecular-mass cationic-active matter*

ISO 2871-2, *Surface active agents — Detergents — Determination of cationic-active matter content — Part 2: Cationic-active matter of low molecular mass (between 200 and 500)*

EAS 794, *Determination of the microbial inhibition of cosmetic soap bars and liquid hand and body washes — Test method*

### 3 Terms and definitions

For the purposes of this document, the terms and definitions given in ISO 862 and the following shall apply.

ISO and IEC maintain terminological databases for use in standardization at the following addresses:

- IEC Electropedia: available at <http://www.electropedia.org/>
- ISO Online browsing platform: available at <http://www.iso.org/obp>

#### 3.1

##### **active matter**

in the formulation, the whole of surface active agents responsible for an activity specified

#### 3.2

##### **anionic surface active**

surface active agents which ionizes in aqueous solution to produce negatively charged organic ions which are responsible for surface active

**3.3 builder**  
complementary component of a detergent usually inorganic, which with reference to the washing action adds its characteristic to those of constituents

**3.4 detergent**  
product specially formulated for cleaning through the process of detergency

**3.5 foam**  
mass of gasses cells separated by thin films of liquid and formed by juxtaposition of bubbles, giving a dispersion in which a large proportion of gas by volume is dispersed in a liquid

**3.6 lot**  
all containers of the same size, shape, and design manufactured from the same raw materials under similar condition of production in a consignment

**3.7 soap**  
an anionic surface active agent which exhibits the phenomenon of reversible hydrolysis by action of water. Their reaction is alkaline

## **4 Requirements**

### **4.1 General requirements**

**4.1.1** 4.1.1 The detergent shall be a clear, homogeneous liquid. It shall remain stable after having been maintained at 5 °C for 24 h and shall remain clear, homogeneous fluid when maintained for a further 24 h at 30 °C.

**4.1.2** The detergent shall be free from visible impurities, and objectionable odour. It shall contain synthetic organic active matter and foam stabilizing agents. It shall be completely miscible with water, and contain no soap.

**4.1.3** Liquid soaps shall be free from visible impurities, and objectionable odour. It shall contain organic active matter. It shall be completely miscible with water

**4.1.4** The active ingredients used shall be biodegradable when tested according to EAS 814

**4.1.5** The detergent shall have good cleaning and lathering properties.

**4.1.5** All the substances used in the liquid detergent shall comply with the requirements of EAS 377 (all parts).

**4.1.6** The soap may contain additives that may boost its performance e.g. sodium bicarbonate; hydrogen peroxide etc

## 4.2 Specific requirements

4.2.1 The detergent shall comply with the specific requirements given in Table 1 when tested in accordance with the test methods specified therein.

**Table 1 — Specific requirements for liquid detergent for household use**

S/No	Characteristic		Requirement		Test method
			General Purpose	Dishwashing	
i.	Matter insoluble in water% (m/m), max		0.5		Annex A
ii.	Rinsing properties		To pass the test		Annex B
iii.	Total active ingredient content, % m/m, min.		5	10	ISO 2870/ISO 2871-1/ ISO 2871-2
iv.	pH at 27 °C of 1 % solution, w/v	products with added weak acids <sup>a</sup>	3.5 <sup>a</sup> - 12		Annex D
		products without added acids	5.0 – 12		
v.	Total non-detergent organic matter, % m/m, max.		0.5		Annex E
vi.	Inorganic salts content, % (m/m), max.		Not applicable	5	Annex F
vii.	Antibacterial properties <sup>b,c</sup>		To pass the test		EAS 794

<sup>a</sup>only weak acids should be used such as lactic acid, citric acid, benzoic acid, salicylic acid, hyaluronic acid, mandelic acid, glycolic acid, etc

<sup>b</sup>only applicable to products claiming antibacterial properties

<sup>c</sup>ASTM E1174 and EN 1276 factoring log reduction may be used as alternative test methods to determine antibacterial activity. When these test methods are used, the log reduction shall not be less than 99%.

4.2.2 The liquid soaps shall comply with the specific requirements given in Table 2 when tested in accordance with the test methods specified therein.

**Table 2— Specific requirements for liquid soaps for household use**

S/No	Characteristic		Requirement		Test method
			General Purpose	Dishwashing	
i.	Total fatty matter, %, min		15		ISO 685
ii.	Rinsing properties		To pass the test		Annex B
iii.	pH at 27 °C of 1 % solution, w/v	With added acids	3.5 <sup>a</sup> – 12		Annex D
		Without added acids	5.0 – 12		
iv.	Matter insoluble in ethanol, % by mass, max		5		ISO 673
v.	Inorganic salts content, % (m/m), max.		Not applicable	5	Annex F
vi.	Antibacterial properties <sup>b,c</sup>		To pass the test		EAS 794

<sup>a</sup>only weak acids should be used such as lactic acid, citric acid, benzoic acid, salicylic acid, hyaluronic acid, mandelic acid, glycolic acid, etc

<sup>b</sup> only applicable to products claiming antibacterial properties

<sup>c</sup>ASTM E1174 and EN 1276 factoring log reduction may be used as alternative test methods to determine antibacterial activity. When these test methods are used, the log reduction shall not be less than 99%.

## 5 Packaging

The product shall be packaged in a suitable, well-closed container to protect the integrity of the product during transportation and storage.

## 6 Labelling

Each container shall be legibly and indelibly labelled in English, and Kiswahili or French or a combination of any other language as agreed to between the manufacturer and supplier with the following information:

- a) name of the product as "liquid detergent" or "liquid soap"
- b) intended use as "general purpose" or "dishwashing"
- c) Clearly indicate whether synthetic, combined or soap-based.
- d) an indication of anti-microbial activity (where applicable)
- e) manufacturer's name and physical address;
- d) batch or code number;
- e) net content;
- f) country of origin;
- g) instructions for use;
- h) date of manufacture"; and

- i) best before date.
- j) list of ingredients

NOTE The name, physical address of the distributor/supplier and trade mark may be added.

## **7 Sampling**

Sampling shall be done in accordance with Annex C

FOR PUBLIC COMMENT ONLY

## Annex A (normative)

### Determination of matter insoluble in water

#### A.1 Principle

A known mass of sample is diluted and filtered. The residues are then dried to constant mass.

#### A.2 Procedure

**A.2.1** Weigh, to the nearest (to  $\pm 0.001$  g) approximately 5 g of the test sample, into a 400-mL beaker and add 200 mL of distilled water. Heat on a steam bath, with frequent stirring, until the sample is completely dispersed.

**A.2.2** Filter the solution immediately, under suction, through a previously dried and tared sintered glass crucible of porosity 2. Ensure that the insoluble matter is quantitatively transferred to the filter.

**A.2.3** Wash the beaker and the residue in the crucible five times with 40 mL of hot distilled water.

**A.2.4** Allow the wash solution to drain completely and dry the crucible to constant mass at  $105\text{ }^{\circ}\text{C} \pm 2\text{ }^{\circ}\text{C}$  in an air oven.

#### A.3 Calculation

The insoluble matter content  $S$ , expressed as percent by mass, shall be calculated as follows:

$$S = \frac{M_4 - M_2}{M_1} \times 100$$

Where,

$M_1$  is the mass, in grams, of the test sample;

$M_2$  is the mass in grams, of the sintered glass crucible; and

$M_4$  is the mass, in grams, of the sintered glass crucible and the residue after drying.

## Annex B (normative)

### Test for rinsing properties

#### B.1 Preparation of synthetic hard water

Weigh to the nearest 0.001 g, about 0.264 g of  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$  and 0.295 g of  $\text{MgSO}_4 \cdot 2\text{H}_2\text{O}$ . Transfer quantitatively to a 1-L volumetric flask; dissolve in a small portion of distilled water and make up to the mark with distilled water. The resulting solution will have a concentration of 8.1 millimole per litre calcium hardness.

#### B.2 Procedure

Dissolve 2.0 mL of the liquid soap or liquid detergent, or combined synthetic and soap as completely as possible in 98 mL of synthetic hard water (see B.1) at ambient temperature, in a clean 250-mL Erlenmeyer flask. Stopper the flask and stir vigorously for 1 min. Pour out the solution. Rinse the flask by the same procedure, using three 75 mL portions of synthetic hard water alone. Invert the flask, allow to dry and examine for any residue not rinsed from the interior. The flask shall contain no more residues after being dried than a similar allowed drying after rinsing with synthetic hard water alone.

## Annex C (normative)

### Sampling

#### C.1 Procedure

G.1.1 In a single consignment, all packages (cartons) containing liquid detergent drawn from the same batch of production shall constitute a lot. For ascertaining the conformity of the lot to the requirements of this standard, tests shall be carried out on each lot separately. The number of packages to be selected for drawing the sample shall be in accordance with Table G.1.

Table G.1 — Scale of sampling

Number of packages (cartons) in the lot N	Number of packages (cartons) to be selected n	Number of samples per carton selected
4 to 15	3	3
6 to 40	4	4
41 to 65	5	2
66 to 110	7	2
111 and above	10	1

**C.1.2** The packages shall be selected at random, using tables of random numbers. If these are not available, the following procedure shall be applied:

Starting from any package, count all the packages in one order as 1, 2, 3... N, selecting every  $k^{\text{th}}$  package, where  $k$  is the integral part of  $N/n$ .

**C.1.3** From each package thus selected, draw at random an equal number of units so as to obtain a total volume of at least 2 L.

#### C.2 Samples for testing

Take at one time all test samples required for the tests in 4.2. Measure the test sample required for determination of free alkali or acid content, and use it immediately.

## Annex D (normative)

### Determination of pH

#### D.1 General

pH determination should be made in an acid free atmosphere.

#### D.2 Apparatus

**D.2.1 Any standard pH meter**, equipped with a low sodium error glass electrode. The instrument shall be calibrated and standardized with standard buffer solutions (see D.3.2) before use.

**D.2.2 Volumetric flask**, 1000-mL capacity

**D.2.3 Beakers**, 1000-mL

#### D.3 Reagents

**D.3.1** Distilled water shall be boiled thoroughly or purged with carbon dioxide-free air to remove carbon dioxide and shall be protected with soda lime or soda asbestos while cooling and in storage. The pH of this water shall be between 6.2 and 7.2 at 27 °C. The residue on evaporation when heated at 105 °C for one hour shall not exceed 0.5 mL/L.

**D.3.2** Standard buffer solutions with the pH range of 9 to 11 at 27 °C for calibrating the pH meter.

#### D.4 Procedure

Weigh to the nearest milligram approximately 10g of the material and transfer to a 1-L volumetric flask. Partially fill the flask with distilled water and agitate until the sample is completely dissolved. Adjust the temperature of the solution and the distilled water to 27 °C ± 2 °C and fill to the calibration mark with distilled water, stopper the flask mix thoroughly and allow the solution to stand at a temperature of 27 °C ± 2 °C for two hours prior to measuring the pH. Measure the pH of the solution at 27 °C ± 2 °C using a glass electrode.

## Annex E (normative)

### Determination of non-detergent organic matter

#### E.1 General

The term non-detergent organic matter includes hydrocarbons, fatty alcohols and perfumes. Using petroleum ether and under the conditions prescribed, non-detergent organic matter only is extracted leaving any alkylamide present in the material.

#### E.2 Apparatus

E.2.1 Evaporating basin

E.2.2 Separating funnels, 1 000-mL capacity.

E.2.3 Wide mouthed flat-bottomed flask, 200-mL capacity

E.2.4 Buchner flask, 500-mL capacity, fitted with a sintered glass filter funnel (porosity 4)

#### E.3 Reagents

E.3.1 Ethyl alcohol, 50 %, 70 %, 90 % and 96% (by volume)

E.3.2 Petroleum ether, boiling range 40 °C to 60 °C non-volatile residue at 80 °C maximum 0.001%

E.3.3 Acetone, non-volatile residue at 80 °C maximum 0.001 %

#### E.4 Procedure

E.4.1 For the removal of inorganic salts, weigh accurately about 5 g of the material in a 150-mL squat beaker. Extract with 50 mL of hot 90 % ethanol by heating on the steam bath for about 2 min stirring and breaking up any hard lumps with a glass rod flattened at the end.

Allow the solid matter to settle and decant the hot alcoholic solution through a sintered glass filter funnel (porosity 4) fitted to a 500-mL Buchner flask to which suction is applied. Repeat the extraction in a similar manner with five further consecutive 30-mL quantities of boiling 90 % ethanol. Pass each extract in turn through the filter into the flask.

E.4.2 Transfer quantitatively all the combined filtrates from the Buchner flask to a 1-L separating funnel and rinse the flask four times with 40-mL quantities of distilled water, transferring each wash in turn to the separating funnel. Add 100 mL of petroleum ether, swirl gently to ensure adequate mixing and allow the two phases to separate. Run off the aqueous alcoholic layer into a second separating funnel, and extract with 75 mL of petroleum ether. Repeat the extraction of the aqueous alcoholic phase in the third separating funnel with a further 75 mL of petroleum ether. Combine the three ether extracts in the first separating funnel. Rinse each of the two empty funnels with a few millilitres petroleum ether and add the rinsing to the combined ether extracts.

E.4.3 Wash the combined ether extracts and rinsing (see E.4.2) with four successive 50-mL portions of 70 % ethyl alcohol, shaking and removing the alcoholic phase each time. Transfer the ether layer in stages to a tared flask and evaporate off the solvent. Add 10 mL of acetone and evaporate off the solvent. Rotate the flask on a steam bath during the operation. Cool the flask to about 60 °C to 65 °C, gently blow out the last traces of solvent with a current of dry air, cool in a desiccator and weigh.

## E.5 Calculation

The non-detergent organic matter, expressed as percent by mass, shall be calculated as follows:

$$100 \frac{m_1}{m}$$

where

$m_1$  is the mass, in grams, of the non-detergent organic matter in the flask; and

$m$  is the mass, in grams, of the material taken for the test.

FOR PUBLIC COMMENT ONLY

## Annex F (normative)

### Determination of inorganic salts

#### F.1 Procedure

Take the dish containing the material after evaporation. Heat it at 450 °C in a muffle furnace to destroy organic matter. Cool the dish and its contents, add a few drops of concentrated sulphuric acid and heat again to dryness. Cool and weigh. Repeat the process of heating, cooling and weighing until constant mass is obtained.

#### F.2 Calculation

The inorganic salts content, expressed as percent by mass, shall be calculated as follows:

$$\left( \frac{M_1 - M_3}{M_1 - M_0} \times 100 \right)$$

where,

$M_0$  is the mass, in grams, of the dish;

$M_1$  is the mass, in grams, of the dish and the sample before heating; and

$M_3$  is the mass, in grams, of the dish and the residue.

## Bibliography

- [1] EAS 383: 2021, *Liquid detergent for household use — Specification*

FOR PUBLIC COMMENT ONLY